



### **Meniscal Repair Post-Operative Instructions**

1. After arthroscopy the wound is covered with gauze or ace wraps. Due to the large amount of fluid used during the arthroscopy, it is normal to see some bloody drainage on the dressings. If bright red blood persists despite elevation and icing, please call the doctor. The dressing should be removed and wounds covered with Band-Aids on the 4th day after surgery. Do not remove the paper strips or cut any of the visible suture. Re-apply the ace wrap for 5-7 days to control swelling. Wounds should be kept dry for 4 days. Showering is allowed with after the 4th day with bandaids over the incisions. The wound should not be submerged in a bathtub or pool until allowed to do so by Dr. Overturf
2. Icing is very important for the first 5-7 days postoperative. While the post-op dressing is in place, icing should be as often as possible. Ice is applied for 20-minute periods 3-4 times per day. Care must be taken with icing to avoid frostbite to the skin.
3. You will be placed in a brace and are to remain non-weight bearing at all times. The brace can be removed for icing while you are sitting or lying but must remain in place during sleep and during ambulation on crutches.
4. The anesthetic drugs used during your surgery may cause nausea for the first 24 hours. If nausea is encountered, drink only clear liquids (i.e. Sprite or 7-up). The only solids should be dry crackers or toast. If nausea and vomiting become severe or the patient shows sign of dehydration (lack of urination) please call the doctor or the surgery center. A low-grade fever (100.5) is not uncommon in the first 24 hours but unusual beyond. Please call the doctor with any temperature over 101.0 degrees.
5. Take a baby aspirin (81 mg) daily until the sutures are removed in the office. This may lower the risk of a blood clot developing after surgery. Should severe calf pain occur or significant swelling of calf and ankle, please call the doctor.
6. Local anesthetics are put into the joint during surgery. It is not uncommon for patients to encounter more pain on the first or second day after surgery. This is the time when swelling peaks. Using the pain medication as directed will help control pain with little risk of complication. Taking pain

medication before bedtime will assist in sleeping. It is important not to drink or drive while taking narcotic medication. If you were prescribed narcotic medication, you can supplement those medications with 200 mg or 400 mg of ibuprofen every 4-6 hours. You should resume your normal medications for other conditions the day after surgery. We have no specific diet restrictions after surgery but extensive use of narcotics can lead to constipation. High fiber diet, lots of fluids, and muscle activity can prevent this occurrence.

7. The doctor will need to reexamine you 7-10 days after knee arthroscopy. Please call the office to schedule a follow-up appointment.